

примененем ультрафонофореза 15 % раствора димефосфона выявлена у больных с субкомпенсированными формами кариеса зубов, патогенетической терапии с сублингвальным применением полиоксидония – у больных с декомпенсированными формами кариеса зубов. Эффективность выразилась в статистически значимой оптимизации характеристик стоматологического статуса по изучаемым клиническим критериям.

CLINICAL EVALUATION OF OPERATIVE RESTORATIVE MANIPULATIONS WITH DIFFERENT FORMS OF DENTAL CARIES

Gilyazeva V.V.

SBEI HPE «Kazan State Medical University» of The Ministry of health of the Russian Federation, (420012, Kazan, Butlerov street, 49) hedstroem@rambler.ru; gilyazevav@mail.ru

The state of fillings / restorations of hard tissue in the area of fillings / restorations in the amount of operative rehabilitation interventions, as well as - as the operating and recovery of manipulation in patients with tooth decay - adolescents aged 15-17 years, who received a variety of methods of pathogenetic therapy in the complex treatment caries, according to clinical criteria Ryge. The results of evaluation of dental status on the studied clinical criteria to identify the different effectiveness of the methods of pathogenetic therapy in sub-and decompensated forms of tooth decay: the greatest efficiency of pathogenetic therapy with phonophoresis 15% solution dimephosphone detected in patients subcompensated form of dental caries, pathogenetic therapy with sublingual application полиоксидония - patients decompensated forms of dental caries. The efficiency has resulted in statistically significant optimization of the characteristics of dental status on the studied clinical criteria.

КЛИНИКО–ЛАБОРАТОРНОЕ ОБОСНОВАНИЕ АЛГОРИТМОВ ПАТОГЕНЕТИЧЕСКОГО ВЕДЕНИЯ БОЛЬНОГО КАРИЕСОМ ЗУБОВ

Гилязева В.В., Гиниятуллин И.И.

ГБОУ ВПО «Казанский государственный медицинский университет» Министерства здравоохранения Российской Федерации, Россия, Татарстан (420012, г. Казань, ул. Бултерова, 49) hedstroem@rambler.ru; gilyazevav@mail.ru

Проведен анализ клинико-лабораторного изучения различных методов патогенетической терапии в составе комплексного лечения больных кариесом зубов – подростков в возрасте 15–17 лет. Результаты исследования позволили установить наибольшую эффективность патогенетической терапии с применением ультрафонофореза 15 % раствора димефосфона и сублингвального применения полиоксидония в составе комплексного лечения больных кариесом зубов при К 02.0, К 02.1 субкомпенсированной и К 02.0, К 02.1 декомпенсированной степени активности патологического процесса. Эта эффективность выразилась в изменении показателей стоматологического статуса – интенсивности и прироста интенсивности кариеса зубов, резистентности эмали, а также – состояния региональной резистентности полости рта при проведении соответствующей патогенетической терапии при изучаемых формах заболевания. Результаты проведенного исследования позволили обосновать и сформулировать основные положения алгоритмов патогенетического ведения больных кариесом зубов в составе комплексного лечения в зависимости от степени активности патологического процесса.

CLINICAL LABORATORY JUSTIFICATION OF ALGORITHMS PATHOGENETIC MANAGING PATIENTS OF DENTAL CARIES

Gilyazeva V.V., Guiniatullin I.I.

SBEI HPE «Kazan State Medical University» of The Ministry of health of the Russian Federation, (420012, Kazan, Butlerov street, 49) hedstroem@rambler.ru; gilyazevav@mail.ru

The analysis of the clinical and laboratory studies of various methods of pathogenetic therapy in the complex treatment of patients with dental caries - adolescents aged 15–17 years. Results of the study to establish the most efficient pathogenetic therapy with phonophoresis 15 % solution dimephosphone and sublingual polyoxidonium in the complex treatment of patients with dental caries to К 02.0, К 02.1 subcompensated and К 02.0, К 02.1 decompensated degree of activity of the pathological process. This efficiency has resulted in a change of dental status indicators – the intensity and increase the intensity of dental caries and enamel resistance and - state of regional resistance during oral therapy with corresponding pathogenic forms of the disease under study. Results of the study allowed to justify and articulate the basic provisions of the algorithms pathogenic case management of dental caries in the complex treatment, depending on the degree of activity of the pathological process.

КОМПЛЕКСНЫЙ ПОДХОД К ОЦЕНКЕ ЭСТЕТИКО-ФУНКЦИОНАЛЬНЫХ РЕЗУЛЬТАТОВ КАРИЕС-ИНФИЛЬТРАЦИИ ПРИ ЛЕЧЕНИИ КАРИЕСА ЭМАЛИ

Гилёва О.С., Муравьёва М.А., Серебренникова Е.В.

Государственное бюджетное образовательное учреждение высшего профессионального образования «Пермская государственная медицинская академия имени академика Е. А. Вагнера» Министерства здравоохранения Российской Федерации», Россия, 614000, Пермь, ул. Петропавловская, 26

В статье обоснована необходимость комплексного подхода к оценке эффективности лечения кариеса эмали методом кариес-инфильтрации, основанного на динамической оценке цвето-текстурных и размерных харак-