

being conducted for the first time in the comparative analysis in the frequency of immunochemical version of kirghiz population and the Russian-speaking population of Kirghizia. According to the results of our study on assessing the impact of immunoglobulin class on outcome of multiple myeloma, the survival rate of patients with Ig A- myeloma, Ig D-myeloma and myeloma IgE, IgA is much lower compared with patients suffering from Ig G-myeloma. MM patients kirghiz nationality recorded in the form of immunochemical embodiment 65 % Ig A–myeloma cases, Ig D-myeloma in 1,5 % cases, IgE myeloma 0,5 % cases, and when detected early uptake of primary chronic kidney disease and myeloma Ig G-recorded in 33 % cases. Based on test results of clinical and laboratory parameters and chronic renal failure in IgA, Ig D, Ig E-myeloma compared with IgG-myeloma for more malignant disease. This requires a timely selection of therapy and chemotherapy program selection.

ИММУНОГЕНЕТИЧЕСКИЕ ПОКАЗАТЕЛИ И РАСПРОСТРАНЕННОСТЬ НОСИТЕЛЬСТВА ВИРУСА ЭПШТЕЙНА - БАРР, ЦИТОМЕГАЛОВИРУСА У ПОТЕНЦИАЛЬНЫХ ДОНОРОВ ГЕМОПОЭТИЧЕСКИХ СТВОЛОВЫХ КЛЕТОК

Баратова М.А.², Баратова Д.А.^{1,3}, Нурлыханов Е.Б.^{1,4}

1 МОО «Национальный Регистр гемопоэтических стволовых клеток Киргизии в Санкт-Петербурге» Санкт-Петербург, Россия. (195197, Санкт-Петербург, проспект Полустровский д.59 литер Ш),

e-mail: baratova@list.ru

2 АО «Республиканский научный центр неотложной медицинской помощи», Астана, Казахстан (010000, Астана, улица Керей-Жанибек хандар д.3),

e-mail: maksat-brt@list.ru

3 НМУ «Евразийский центр онкогематологии, иммунологии и терапии», Санкт-Петербург, Россия (195197, Санкт-Петербург, проспект Полустровский д. 59 литер Ш),

e-mail: baratova@list.ru

4 ГУ «Областная клиническая больница» Шымкент, Казахстан (160011, Шымкент, улица Майлы-Кожа д. 4), e-mail: tavshan7@mail.ru

В статье представлены результаты исследований иммуногенетических и популяционных характеристик доноров костного мозга киргизской национальности в сравнении со здоровыми жителями Северо-западного региона Российской Федерации из Российского регистра костного мозга. Установлены высоко достоверные отличия, выявлены значимые HLA-маркеры у киргизской нации; при исследовании на вирусы в группе доноров выявлен вирус Эпштейна-Барр в 26 % и цитомегаловирус в 39 % случаях. У потенциальных доноров гемопоэтических стволовых клеток с хроническими очагами инфекций выявлены повышенные титры иммуноглобулинов IgA k, l и Ig G k, l легкими цепями. Таким образом, выявленные различия в сопоставлении с контрольной популяционной группой ставят вопрос о своевременном подборе и выборе потенциальных доноров костного мозга. При наличии HLA-идентичного донора применение аллогенной трансплантации гемопоэтических стволовых клеток. Для качественного подбора донора - улучшить алгоритм безопасности донора костного мозга. На современном этапе трансплантация костного мозга дает наилучшие результаты для увеличения показателей выживаемости при лечении злокачественных заболеваний.

IMMUNOGENETICHESKIE INDICATORS AND FREQUENCY OF THE CARRIER EPCHTEINA- BARR VIRUS, CYTOMEGALOVIRUS AMONG POTENTIAL DONORS OF HEMOPOETIC STEM CELLS

Baratova M.A.², Baratova D.A.^{1,3}, Nurlyhanov E.B.^{1,4}

1 National Register of hematopoietic stem cells Kirghizia in Saint Petersburg, Saint Petersburg, Russia (195197, St. Petersburg, prospect Polyustrovsky 59, literU),

e-mail: baratova@list.ru

2 Republican Scientific Center of Emergency Medical Assistance, Astana, Kazakhstan (010000, Astana, street-Kerey-Janibek handar, 3),

e-mail: maksat-brt@list.ru

3 Eurasian Center oncohematology, immunology and therapy, Saint Petersburg, Russia (195197, St.Petersburg, prospect Polyustrovsky 59,literU),

e-mail: baratova@list.ru

4 Regional Clinical Hospital” Shymkent, Kazakhstan, (160011, Shymkent, street Maily -Kojo, 4), e-mail:tavshan7@mail.ru

The article presents the results of studies of immunogenetic and population characteristics of bone marrow donors in the kyrgyz ethnic group compared with healthy residents of the Northwest region of the Russian Federation of the Russian register of bone marrow. Set high significant differences were revealed significant HLA-markers in the kyrgyz nation; viruses when tested on a group of donors identified Epstein - Barr virus in 26% and 39% cytomegalovirus cases. Among potential donors of hematopoietic stem cells with chronic foci of infection revealed increased titers of immunoglobulin Ig A k, l and Ig G k,l, light chains. Thus, the observed differences in comparison with the control group raises the question of a timely selection of treatment and choice of potential bone marrow donors. At presence of HLA-identical donor use of allogeneic hematopoietic stem cell transplantation. For qualitative selection of the donor - to improve the security algorithm and a bone marrow donor. At the present stage, bone marrow transplantation provides the best results for increasing survival in the treatment of malignant diseases.